ELSEVIER

Carbohydrate Research 286 (1996) C7–C8

CARBOHYDRATE RESEARCH

Contents

Regular Papers

NMR Study of uronic acids and their complexation with molybdenum(VI) and tungsten(VI) oxoions
M.L.D. Ramos, M.M.M. Caldeira and V.M.S. Gil (Coimbra, Portugal) 1

Binding properties of carbohydrate sulfamates based on ab initio 6-31 + G** calculations on *N*-methyl and *N*-ethyl sulfamate anions
D.M. Whitfield (Ottawa, Canada), D. Lamba (Rome, Italy), T.-H. Tang (Toronto, Canada; Tianjin, People's Republic of China) and I.G. Csizmadia (Toronto, Canada) 17

Quantitative determination of native and methylated cyclodextrins by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry
H. Bartsch, W.A. König, M. Straßner and U. Hintze (Hamburg, Germany) 41

Influence of intramolecular hydrogen-bonding on the conformation of 3-deoxy-3-thioureido sugars
J.M. García Fernández, C. Ortiz Mellet, J.L. Jiménez Blanco, J. Fuentes, M. Jesús Diánez, M.D. Estrada, A. López-Castro and S. Pérez-Garrido (Seville, Spain) 55

A multinuclear NMR spectroscopy study of the tungstate and molybdate complexes of D-fructose and L-sorbose
J.-P. Sauvage, J.-F. Verchère and S. Chapelle (Mont-Saint-Aignan, France). 67

A novel approach to the synthesis of lipophilic thymidinemonophosphoglucopyranosides as drug delivery systems
A. De Nino, A. Liguori, A. Procopio, E. Roberti and G. Sindona (Arcavacata di Rende, Italy). . . . 77

Recognition of β-D-Gal *p*-(1 → 3)-β-D-Glc *p*NAc-OR acceptor analogues by the Lewis α-(1 → 3/4)-fucosyltransferase from human milk
M. Du and O. Hindsgaul (Edmonton, Canada) 87

Synthesis of kanamycin A analogs containing 6-amino-3-oxa-2,3,4,6-tetradeoxy-D- and -L-*glycero*-hexopyranose
R. Kuwahara and T. Tsuchiya (Kawasaki, Japan). 107

Synthesis of novel vitamin C phosphodiesters: Stability and antioxidant activity
K. Morisaki and S. Ozaki (Matsuyama, Japan) 123

Degradation of pectins in alkaline conditions: kinetics of demethylation
C.M.G.C. Renard and J.-F. Thibault (Nantes, France) 139

Notes

A detailed ^{1}H and ^{13}C NMR study of a repeating disaccharide of hyaluronan: the effect of sodium and calcium ions
W. Sicińska and L.E. Lerner (Madison, WI, USA) 151

Synthesis of 1,2-anhydro-3,5,6-tri-*O*-benzyl-α-L-gulofuranose and 1,2-anhydro-3,5-di-*O*-benzyl-β-D-arabinofuranose
X. Ding and F. Kong (Beijing, People's Republic of China) 161

Preparation of amidine derivatives from 1-cyano-D-galactal
L. Somsák (Debrecen, Hungary) 167

A practical synthesis from siastatin B of (3*S*,4*S*,5*R*,6*R*)-4,5-dihydroxy-6-(trifluoroacetamido)piperidine-3-carboxylic acid having antimetastatic activity in mice
T. Satoh, Y. Nishimura, S. Kondo and T. Takeuchi (Tokyo, Japan) 173

Identification of per-*O*-(trimethylsilyl) derivatives of aldoses generated from thermal decomposition of *N*-(1-deoxy-D-fructopyranos-1-yl)proline: reversibility of the Amadori rearrangement
V.A. Yaylayan and A. Huyghues-Despointes (Ste. Anne de Bellevue, Canada) 179

Mixed-acetal glucopyranosides of 2-bromo- and 2-iodo-acetaldehyde by mild alkylation of lactol groups. Irreversible cross inhibition using a pair of glycosidases
H. Ebrahim, D.J. Evans (London, UK), J. Lehmann and L. Ziser (Freiburg, Germany) 189

Author index C1

Subject index C3

Contents C7